

A Phase 3 Randomized Study Comparing Talquetamab SC in Combination With Daratumumab SC and Pomalidomide (Tal-DP) or Talquetamab SC in Combination With Daratumumab SC (Tal-D) Versus Daratumumab SC, Pomalidomide and Dexamethasone (DPd), in Participants With Relapsed or Refractory Multiple Myeloma who Have Received at Least 1 Prior Line of Therapy

Estado Fin Reclutamiento	Tipo de Participantes No aportado	Rangos de Edad Mayores de 64 , Adultos (18-64 años)
Género Ambos	Fases Fase III	Participantes esperados 553
Resultados Sin resultados	Bajo nivel intervención No	Enfermedad rara Si
Cobertura geográfica Sin determinar	Ámbitos del ensayo tratamiento, seguridad, eficacia, farmacocinética, farmacodinámica	Terapia avanzada NO

Información

Identificador
2023-503467-41-00 (CTIS)

Cod. Protocolo
64407564MMY3002

Transicionado 2021-000202-22

Área terapéutica
Enfermedades [C] - Cáncer [C04]

Enfermedad investigada
Relapsed or Refractory Multiple Myeloma

Título Científico

A Phase 3 Randomized Study Comparing Talquetamab SC in Combination With Daratumumab SC and Pomalidomide (Tal-DP) or Talquetamab SC in Combination With Daratumumab SC (Tal-D) Versus Daratumumab SC, Pomalidomide and Dexamethasone (DPd), in Participants With Relapsed or Refractory Multiple Myeloma who Have Received at Least 1 Prior Line of Therapy

Justificación

No aportado

Objetivo Principal

To compare the efficacy of Tal-DP and Tal-D, respectively, with DPd

Variables de Evaluación Primaria

Progression-Free Survival

Momentos temporales de evaluación primaria

No aportado

Objetivo Secundario

No aportado

Variables de Evaluación Secundaria

No aportado

Momentos temporales de evaluación secundaria

No aportado

Criterios de Inclusión

Documented multiple myeloma as defined: a) Multiple myeloma diagnosis according to the International Myeloma Working Group (IMWG) diagnostic criteria and b) Measurable disease at screening as defined by any of the following: i) Serum M-protein level greater than or equal to ($\geq$) 0.5 grams per deciliter (g/dL) (central laboratory); ii) Urine M-protein level $\geq$ 200 milligram (mg) per 24 hours (central laboratory); iii) Light chain multiple myeloma

without measurable M-protein in the serum or the urine: serum immunoglobulin free light chain ≥ 10 milligram per deciliter (mg/dL) (central laboratory), and abnormal serum immunoglobulin kappa lambda free light chain ratio Relapsed or refractory disease as defined by: i) Relapsed disease is defined as an initial response to prior treatment, followed by confirmed progressive disease by IMWG criteria greater than ($>$) 60 days after cessation of treatment; ii) Refractory disease is defined as less than ($<$) 25 percent (%) reduction in monoclonal paraprotein (M-protein) or confirmed progressive disease by IMWG criteria during previous treatment or less than or equal to ($\leq$) 60 days after cessation of treatment Received at least 1 prior line of antimyeloma therapy including a proteasome inhibitor (PI) and lenalidomide. Participants who have received only 1 prior line of antimyeloma therapy must be considered lenalidomide-refractory (that is, have demonstrated progressive disease by IMWG criteria on or within 60 days of completion of lenalidomide-containing regimen). Participants who have received ≥ 2 prior lines of antimyeloma therapy must be considered lenalidomide exposed Documented evidence of progressive disease based on investigator's determination of response by the IMWG criteria on or after their last regimen Have an Eastern Cooperative Oncology Group (ECOG) performance status score of 0, 1, or 2 at screening and immediately prior to the start of administration of study treatment

Criterios de Exclusión

Contraindications or life-threatening allergies, hypersensitivity, or intolerance to study drug excipients Disease is considered refractory to an anti-cluster of differentiation 38 (CD38) monoclonal antibody as defined per IMWG consensus guidelines (progression during treatment or within 60 days of completing therapy with an anti-CD38 monoclonal antibody) Received prior pomalidomide therapy A maximum cumulative dose of corticosteroids to ≥ 140 milligrams (mg) of prednisone or equivalent within 14-day period before the first dose of study drug Known active central nervous system (CNS) involvement or exhibits clinical signs of meningeal involvement of multiple myeloma. If either is suspected, negative whole brain magnetic resonance imaging (MRI) and lumbar cytology are required Plasma cell leukemia (per IMWG criteria) at the time of screening, Waldenström's macroglobulinemia, polyneuropathy, organomegaly, endocrinopathy, M-protein, and skin changes (POEMS syndrome), or primary amyloid light chain amyloidosis

Calendario

(Última actualización: 23/01/2026)

Autorización 26/10/2022	Inicio de Ensayo 03/11/2022	Inclusión Primer Paciente 08/11/2022	Interrumpido No aportado	Reiniciado No aportado
Fin de reclutamiento 19/03/2024	Fin prematuro (España) No aportado	Fin prematuro (Global) No aportado	Fin del ensayo en España No aportado	Fin del ensayo global No aportado

Promotor

Janssen - Cilag International Belgium

Turnhoutseweg 30

Contact Person

CTIS Point of Contact

+31717996133

CTISadmin@its.jnj.com

Monetary support: N/A

Tipo de promotor: Comercial

Ceim

CEIm de la Comunidad Foral de Navarra

C/ Irunlarrea, 3 31008 Pamplona

ceic@cfnavarra.es

Centros

No iniciado (---/---/---)	Clinica Universidad De Navarra Pamplona NAVARRA Servicio de Hematología
Activo (29/08/2023)	Clínica Universidad De Navarra Pamplona/Iruña NAVARRA
Activo (15/11/2022)	Complejo Asistencial Universitario De Salamanca Salamanca SALAMANCA
Activo (12/12/2022)	Complejo Hospitalario Universitario De Canarias San Cristóbal de La Laguna STA. CRUZ DE TENERIFE
No iniciado (---/---/---)	Fundacio Institut De Recerca De L Hospital De La Santa Creu I Sant Pau Barcelona BARCELONA Servicio de Hematología
Activo (28/11/2022)	HOSPITAL CLINIC DE BARCELONA Barcelona BARCELONA
No iniciado (---/---/---)	Hospital Clinic De Barcelona Barcelona BARCELONA Servicio de Hematología
Activo (22/12/2022)	HOSPITAL CLINICO UNIVERSITARIO VIRGEN DE LA ARRIXACA Murcia MURCIA

No iniciado (--)

Hospital De Jerez De La Frontera

Jerez De La Frontera

CÁDIZ

Servicio de Hematología

Activo (03/11/2022)

HOSPITAL DE LA SANTA CREU I SANT PAU

Barcelona

BARCELONA

Activo (19/01/2023)

HOSPITAL DE LEON (COMPLEJO ASISTENCIAL UNIVERSITARIO DE LEON)

León

LEÓN

No iniciado (--)

Hospital Germans Trias I Pujol

Badalona

BARCELONA

Servicio de Hematología

Activo (09/03/2023)

HOSPITAL UNIVERSITARI GERMANS TRIAS I PUJOL DE BADALONA

Badalona

BARCELONA

No iniciado (--)

Hospital Universitari Vall D Hebron

Barcelona

BARCELONA

Servicio de Hematología

Activo (23/11/2022)

HOSPITAL UNIVERSITARI VALL D'HEBRON

Barcelona

BARCELONA

No iniciado (--)

Hospital Universitario De Canarias

La Laguna

STA. CRUZ DE TENERIFE

Servicio de Hematología

Activo (03/11/2022)

HOSPITAL UNIVERSITARIO DE JEREZ DE LA FRONTERA

Jerez de la Frontera

CÁDIZ

Cerrado (16/05/2024)

HOSPITAL UNIVERSITARIO DE LA PRINCESA

Madrid

MADRID

No iniciado (---/---/----)

Hospital Universitario De Leon

Leon

LEÓN

Servicio de Hematología

No iniciado (---/---/----)

Hospital Universitario De Salamanca

Salamanca

SALAMANCA

Servicio de Hematología

Activo (25/11/2022)

HOSPITAL UNIVERSITARIO DR. PESET ALEIXANDRE

València

VALENCIA

No iniciado (---/---/----)

Hospital Universitario Dr Peset Aleixandre

Valencia

VALENCIA

Servicio de Hematología

Cerrado (29/01/2025)

Hospital Universitario Hm Sanchinarro

Madrid

MADRID

Cerrado (16/05/2024)

Hospital Universitario Ramón Y Cajal

Madrid

MADRID

Activo (16/12/2022)

HOSPITAL UNIVERSITARIO VIRGEN DE LA VICTORIA

Málaga

MÁLAGA

No iniciado (---/---/----)

Hospital Universitario Virgen De La Victoria

Malaga

MÁLAGA

Servicio de Hematología

Activo (18/11/2022)

HOSPITAL UNIVERSITARIO 12 DE OCTUBRE

Madrid

MADRID

No iniciado (---/---/----)

Hospital Universitario 12 De Octubre

Madrid

MADRID

Servicio de Hematología

No iniciado (---/---/----)

University Clinical Hospital Virgen De La Arrixaca

Murcia

MURCIA

Servicio de Hematología

Medicamentos

JNJ-64407564

SOLUTION FOR INJECTION
SUBCUTANEOUS USE

—

Principios Activos: N/A

Huérfano

Experimental

Pomalidomide

CAPSULE, HARD
ORAL USE

—

Principios Activos: N/A

Experimental

Privigen 100 mg/ml solution for infusion

SOLUTION FOR INFUSION
INTRAVENOUS USE

Código ATC: J06BA02 - INMUNOGLOBULINAS HUMANAS
NORMALES PARA ADM. INTRAVASCULAR

Principios Activos: N/A

Experimental

Privigen 100 mg/ml solution for infusion

SOLUTION FOR INFUSION
INTRAVENOUS USE

Código ATC: J06BA02 - INMUNOGLOBULINAS HUMANAS
NORMALES PARA ADM. INTRAVASCULAR

Principios Activos: N/A

Experimental

Dexamethason 4 mg JENAPHARM®;

TABLET
ORAL USE

Código ATC: H02AB02 - DEXAMETASONA

Principios Activos: N/A

Experimental

KIOVIG 100 mg/ml solution for infusion

SOLUTION FOR INFUSION
INTRAVENOUS USE

Código ATC: J06BA02 - INMUNOGLOBULINAS HUMANAS
NORMALES PARA ADM. INTRAVASCULAR

Principios Activos: N/A

Experimental

Imnovid 2 mg hard capsules

CAPSULE, HARD
ORAL USE

Código ATC: L04AX06 - POMALIDOMIDA

Principios Activos: N/A

Experimental

Pomalidomide

CAPSULE, HARD
ORAL USE

—

Principios Activos: N/A

Experimental

Fortecortin®; 2 mg Tabletten

TABLET
ORAL USE

Código ATC: H02AB02 - DEXAMETASONA
Principios Activos: N/A

Experimental

Dexamethasone Tablets BP 2.0mg

TABLET
ORAL USE

Código ATC: H02AB02 - DEXAMETASONA
Principios Activos: N/A

Experimental

Pomalidomide

CAPSULE, HARD
ORAL USE

–
Principios Activos: N/A

Experimental

KIOVIG 100 mg/ml solution for infusion

SOLUTION FOR INFUSION
INTRAVENOUS USE

Código ATC: J06BA02 - INMUNOGLOBULINAS HUMANAS
NORMALES PARA ADM. INTRAVASCULAR
Principios Activos: N/A

Experimental

Imnovid 4 mg hard capsules

CAPSULE, HARD
ORAL USE

Código ATC: L04AX06 - POMALIDOMIDA
Principios Activos: N/A

Experimental

Imnovid 1 mg hard capsules

CAPSULE, HARD
ORAL USE

Código ATC: L04AX06 - POMALIDOMIDA
Principios Activos: N/A

Experimental

Imnovid 3 mg hard capsules

CAPSULE, HARD
ORAL USE

KIOVIG 100 mg/ml solution for infusion

SOLUTION FOR INFUSION
INTRAVENOUS USE

Código ATC: L04AX06 - POMALIDOMIDA
Principios Activos: N/A

Experimental

Código ATC: J06BA02 - INMUNOGLOBULINAS HUMANAS
NORMALES PARA ADM. INTRAVASCULAR
Principios Activos: N/A

Experimental

Privigen 100 mg/ml solution for infusion
SOLUTION FOR INFUSION
INTRAVENOUS USE

Código ATC: J06BA02 - INMUNOGLOBULINAS HUMANAS
NORMALES PARA ADM. INTRAVASCULAR
Principios Activos: N/A

Experimental

JNJ-64407564
SOLUTION FOR INJECTION
SUBCUTANEOUS USE

-
Principios Activos: N/A

Huérfano

Experimental

DARZALEX 1800 mg solution for injection
SOLUTION FOR INJECTION
SUBCUTANEOUS USE

Código ATC: L01FC01 - DARATUMUMAB
Principios Activos: N/A

Huérfano

Experimental

Pomalidomide
CAPSULE, HARD
ORAL USE

-
Principios Activos: N/A

Experimental

Sin resultados

A Phase 3 Randomized Study Comparing Talquetamab SC in Combination With Daratumumab SC and Pomalidomide (Tal-DP) or Talquetamab SC in Combination With Daratumumab SC (Tal-D) Versus Daratumumab SC, Pomalidomide and Dexamethasone (DPd), in Participants With Relapsed or Refractory Multiple Myeloma who Have Received at Least 1 Prior Line of Therapy

State
End of recruitment

Type of participants
Not provided

Age Ranges
Older than 64 , Adults (18-64 years)

Gender
Both

Phases
Phase III

Expected Participants
553

Results
No results

Low level of intervention
No

Rare disease
Yes

Geographic coverage
Undetermined

Areas of the study
treatment, safety, effectiveness,
pharmacokinetics, pharmacodynamics

Advanced therapy
NO

Information

Identifier
2023-503467-41-00 (CTIS)

Protocol Code
64407564MMY3002

Transitioned 2021-000202-22

Therapeutic area
Diseases [C] - Cancer [C04]

Investigated Disease
Relapsed or Refractory Multiple Myeloma

Scientific Title

A Phase 3 Randomized Study Comparing Talquetamab SC in Combination With Daratumumab SC and Pomalidomide (Tal-DP) or Talquetamab SC in Combination With Daratumumab SC (Tal-D) Versus Daratumumab SC, Pomalidomide and Dexamethasone (DPd), in Participants With Relapsed or Refractory Multiple Myeloma who Have Received at Least 1 Prior Line of Therapy

Rationale

Not provided

Main Objective

To compare the efficacy of Tal-DP and Tal-D, respectively, with DPd

Primary Endpoints

Progression-Free Survival

Temporary moments of secondary assessment

Not provided

Secondary Objective

Not provided

Secondary Endpoints

Not provided

Temporary moments of secondary assessment

Not provided

Inclusion criteria

Documented multiple myeloma as defined: a) Multiple myeloma diagnosis according to the International Myeloma Working Group (IMWG) diagnostic criteria and b) Measurable disease at screening as defined by any of the following: i) Serum M-protein level greater than or equal to ($\geq$) 0.5 grams per deciliter (g/dL) (central laboratory); ii) Urine M-protein level $\geq$ 200 milligram (mg) per 24 hours (central laboratory); iii) Light chain multiple myeloma

without measurable M-protein in the serum or the urine: serum immunoglobulin free light chain ≥ 10 milligram per deciliter (mg/dL) (central laboratory), and abnormal serum immunoglobulin kappa lambda free light chain ratio

Relapsed or refractory disease as defined by: i) Relapsed disease is defined as an initial response to prior treatment, followed by confirmed progressive disease by IMWG criteria greater than ($>$) 60 days after cessation of treatment; ii) Refractory disease is defined as less than ($<$) 25 percent (%) reduction in monoclonal paraprotein (M-protein) or confirmed progressive disease by IMWG criteria during previous treatment or less than or equal to ($\leq$) 60 days after cessation of treatment

Received at least 1 prior line of antimyeloma therapy including a proteasome inhibitor (PI) and lenalidomide. Participants who have received only 1 prior line of antimyeloma therapy must be considered lenalidomide-refractory (that is, have demonstrated progressive disease by IMWG criteria on or within 60 days of completion of lenalidomide-containing regimen). Participants who have received ≥ 2 prior lines of antimyeloma therapy must be considered lenalidomide exposed

Documented evidence of progressive disease based on investigator's determination of response by the IMWG criteria on or after their last regimen

Have an Eastern Cooperative Oncology Group (ECOG) performance status score of 0, 1, or 2 at screening and immediately prior to the start of administration of study treatment

Exclusion criteria

Contraindications or life-threatening allergies, hypersensitivity, or intolerance to study drug excipients

Disease is considered refractory to an anti-cluster of differentiation 38 (CD38) monoclonal antibody as defined per IMWG consensus guidelines (progression during treatment or within 60 days of completing therapy with an anti-CD38 monoclonal antibody)

Received prior pomalidomide therapy

A maximum cumulative dose of corticosteroids to ≥ 140 milligrams (mg) of prednisone or equivalent within 14-day period before the first dose of study drug

Known active central nervous system (CNS) involvement or exhibits clinical signs of meningeal involvement of multiple myeloma. If either is suspected, negative whole brain magnetic resonance imaging (MRI) and lumbar cytology are required

Plasma cell leukemia (per IMWG criteria) at the time of screening, Waldenström's macroglobulinemia, polyneuropathy, organomegaly, endocrinopathy, M-protein, and skin changes (POEMS syndrome), or primary amyloid light chain amyloidosis

Calendar

(Last Update: 23/01/2026)

Authorization 26/10/2022	Start of Trial 03/11/2022	First patient inclusion 08/11/2022	Halted Not aported	Restarted Not aported
------------------------------------	-------------------------------------	--	------------------------------	---------------------------------

End of recruitment 19/03/2024	Premature end (Spain) Not aported	Premature End (Global) Not aported	Trial end (Spain) Not aported	Trial end (Global) Not aported
---	---	--	---	--

Sponsor

Janssen - Cilag International Belgium

Turnhoutseweg 30

Contact Person

CTIS Point of Contact

+31717996133

CTISadmin@its.jnj.com

Monetary support: N/A

Sponsor type: Commercial

Ceim

CEIm de la Comunidad Foral de Navarra

C/ Irunlarrea, 3 31008 Pamplona

ceic@cfnavarra.es

Sites

not initialized (---/---/----)	Clinica Universidad De Navarra Pamplona NAVARRA Servicio de Hematología	Clínica Universidad De Navarra Pamplona/Iruña NAVARRA Active (29/08/2023)
Active (15/11/2022)	Complejo Asistencial Universitario De Salamanca Salamanca SALAMANCA	Complejo Hospitalario Universitario De Canarias San Cristóbal de La Laguna STA. CRUZ DE TENERIFE Active (12/12/2022)
not initialized (---/---/----)	Fundacio Institut De Recerca De L Hospital De La Santa Creu I Sant Pau Barcelona BARCELONA Servicio de Hematología	HOSPITAL CLINIC DE BARCELONA Barcelona BARCELONA Active (28/11/2022)
not initialized (---/---/----)	Hospital Clinic De Barcelona Barcelona BARCELONA Servicio de Hematología	HOSPITAL CLINICO UNIVERSITARIO VIRGEN DE LA ARRIXACA Murcia MURCIA Active (22/12/2022)

not initialized (---/---/----)

Hospital De Jerez De La Frontera

Jerez De La Frontera

CÁDIZ

Servicio de Hematología

Active (03/11/2022)

HOSPITAL DE LA SANTA CREU I SANT PAU

Barcelona

BARCELONA

Active (19/01/2023)

HOSPITAL DE LEON (COMPLEJO ASISTENCIAL UNIVERSITARIO DE LEON)

León

LEÓN

not initialized (---/---/----)

Hospital Germans Trias I Pujol

Badalona

BARCELONA

Servicio de Hematología

Active (09/03/2023)

HOSPITAL UNIVERSITARI GERMANS TRIAS I PUJOL DE BADALONA

Badalona

BARCELONA

not initialized (---/---/----)

Hospital Universitari Vall D Hebron

Barcelona

BARCELONA

Servicio de Hematología

Active (23/11/2022)

HOSPITAL UNIVERSITARI VALL D'HEBRON

Barcelona

BARCELONA

not initialized (---/---/----)

Hospital Universitario De Canarias

La Laguna

STA. CRUZ DE TENERIFE

Servicio de Hematología

Active (03/11/2022)

HOSPITAL UNIVERSITARIO DE JEREZ DE LA FRONTERA

Jerez de la Frontera

CÁDIZ

Closed (16/05/2024)

HOSPITAL UNIVERSITARIO DE LA PRINCESA

Madrid

MADRID

not initialized (---/---/----)

Hospital Universitario De Leon

Leon

LEÓN

Servicio de Hematología

not initialized (---/---/----)

Hospital Universitario De Salamanca

Salamanca

SALAMANCA

Servicio de Hematología

Active (25/11/2022)

HOSPITAL UNIVERSITARIO DR. PESET ALEIXANDRE

València

VALENCIA

not initialized (---/---/----)

Hospital Universitario Dr Peset Aleixandre

Valencia

VALENCIA

Servicio de Hematología

Closed (29/01/2025)

Hospital Universitario Hm Sanchinarro

Madrid

MADRID

Closed (16/05/2024)

Hospital Universitario Ramón Y Cajal

Madrid

MADRID

Active (16/12/2022)

HOSPITAL UNIVERSITARIO VIRGEN DE LA VICTORIA

Málaga

MÁLAGA

not initialized (---/---/----)

Hospital Universitario Virgen De La Victoria

Malaga

MÁLAGA

Servicio de Hematología

Active (18/11/2022)

HOSPITAL UNIVERSITARIO 12 DE OCTUBRE

Madrid

MADRID

not initialized (---/---/----)

Hospital Universitario 12 De Octubre

Madrid

MADRID

Servicio de Hematología

not initialized (---/---/----)

University Clinical Hospital Virgen De La Arrixaca

Murcia

MURCIA

Servicio de Hematología

Medication

JNJ-64407564

SOLUTION FOR INJECTION
SUBCUTANEOUS USE

–

Active Principles: N/A

Orphan

Experimental

Pomalidomide

CAPSULE, HARD
ORAL USE

–

Active Principles: N/A

Experimental

Privigen 100 mg/ml solution for infusion

SOLUTION FOR INFUSION
INTRAVENOUS USE

ATC code: J06BA02 - INMUNOGLOBULINAS HUMANAS NORMALES
PARA ADM. INTRAVASCULAR

Active Principles: N/A

Experimental

Privigen 100 mg/ml solution for infusion

SOLUTION FOR INFUSION
INTRAVENOUS USE

ATC code: J06BA02 - INMUNOGLOBULINAS HUMANAS NORMALES
PARA ADM. INTRAVASCULAR

Active Principles: N/A

Experimental

Dexamethason 4 mg JENAPHARM®;

TABLET
ORAL USE

ATC code: H02AB02 - DEXAMETASONA

Active Principles: N/A

Experimental

KIOVIG 100 mg/ml solution for infusion

SOLUTION FOR INFUSION
INTRAVENOUS USE

ATC code: J06BA02 - INMUNOGLOBULINAS HUMANAS NORMALES
PARA ADM. INTRAVASCULAR

Active Principles: N/A

Experimental

Imnovid 2 mg hard capsules

CAPSULE, HARD
ORAL USE

ATC code: L04AX06 - POMALIDOMIDA

Active Principles: N/A

Experimental

Pomalidomide

CAPSULE, HARD
ORAL USE

–

Active Principles: N/A

Experimental

Fortecortin®; 2 mg Tabletten

TABLET

ORAL USE

ATC code: H02AB02 - DEXAMETASONA

Active Principles: N/A

Experimental

Dexamethasone Tablets BP 2.0mg

TABLET

ORAL USE

ATC code: H02AB02 - DEXAMETASONA

Active Principles: N/A

Experimental

Pomalidomide

CAPSULE, HARD

ORAL USE

-

Active Principles: N/A

Experimental

KIOVIG 100 mg/ml solution for infusion

SOLUTION FOR INFUSION

INTRAVENOUS USE

ATC code: J06BA02 - INMUNOGLOBULINAS HUMANAS NORMALES PARA ADM. INTRAVASCULAR

Active Principles: N/A

Experimental

Imnovid 4 mg hard capsules

CAPSULE, HARD

ORAL USE

ATC code: L04AX06 - POMALIDOMIDA

Active Principles: N/A

Experimental

Imnovid 1 mg hard capsules

CAPSULE, HARD

ORAL USE

ATC code: L04AX06 - POMALIDOMIDA

Active Principles: N/A

Experimental

Imnovid 3 mg hard capsules

CAPSULE, HARD

ORAL USE

KIOVIG 100 mg/ml solution for infusion

SOLUTION FOR INFUSION

INTRAVENOUS USE

ATC code: L04AX06 - POMALIDOMIDA

Active Principles: N/A

Experimental

ATC code: J06BA02 - INMUNOGLOBULINAS HUMANAS NORMALES PARA ADM. INTRAVASCULAR

Active Principles: N/A

Experimental

Privigen 100 mg/ml solution for infusion

SOLUTION FOR INFUSION

INTRAVENOUS USE

ATC code: J06BA02 - INMUNOGLOBULINAS HUMANAS NORMALES PARA ADM. INTRAVASCULAR

Active Principles: N/A

Experimental

JNJ-64407564

SOLUTION FOR INJECTION

SUBCUTANEOUS USE

-

Active Principles: N/A

Orphan

Experimental

DARZALEX 1800 mg solution for injection

SOLUTION FOR INJECTION

SUBCUTANEOUS USE

ATC code: L01FC01 - DARATUMUMAB

Active Principles: N/A

Orphan

Experimental

Pomalidomide

CAPSULE, HARD

ORAL USE

-

Active Principles: N/A

Experimental

No results