

A trial to learn how well surovatamig (AZD0486) works and how safe it is in adults with certain blood cancers

Estado Reclutando	Tipo de Participantes No aportado	Rangos de Edad Mayores de 64 , Adultos (18-64 años)
Género Ambos	Fases Fase II	Participantes esperados 188
Resultados Sin resultados	Bajo nivel intervención No	Enfermedad rara No
Cobertura geográfica Sin determinar	Ámbitos del ensayo tratamiento, seguridad, eficacia, farmacocinética, farmacodinamica, farmacogenética, farmacogenómica	Terapia avanzada NO

Información

Identificador 2023-505789-27-00 (CTIS)	Cod. Protocolo No aportado
--	--------------------------------------

Área terapéutica
Enfermedades [C] - Hematología [C15]

Enfermedad investigada
Relapsed or Refractory Non-Hodgkin Lymphoma

Título Científico
A Modular Phase II, Single-arm, Multicenter, Open-label Study to Evaluate the Efficacy and Safety of Surovatamig (AZD0486) in Participants with Relapsed or Refractory B-cell Non-Hodgkin Lymphoma (SOUNDTRACK-B))

Justificación

No aportado

Objetivo Principal

To evaluate the efficacy of surovatamig (AZD0486) single therapy administered at the Recommended Phase 2 Dose (RP2D) in relapsed or refractory B-cell NHL

Variables de Evaluación Primaria

ORR, defined as the proportion of participants achieving either a PR or CR based on Lugano 2014 Response Criteria as determined by an IRC

Momentos temporales de evaluación primaria

No aportado

Objetivo Secundario

1. To assess the safety and tolerability of surovatamig
 2. To further evaluate the efficacy of surovatamig by assessment of DoR and CR rate
 3. To evaluate ORR and CR rate
 4. To evaluate the efficacy of surovatamig by evaluation of additional efficacy measures based on Lugano 2014 Response Criteria
 5. To characterize the PK of surovatamig
 6. To describe the incidence of immunogenicity
 7. To evaluate patient-reported tolerability of surovatamig, including severity of key treatment-related symptoms and overall side-effect burden
 8. To evaluate patient-reported severity of key disease-related symptoms, as well as the impact of disease on lymphoma-specific concerns and general HRQoL, while on surovatamig
 9. Evaluate the impact of surovatamig on MRD-negative CR rate
-

Variables de Evaluación Secundaria

1. Safety and tolerability of surovatamig - incidence, nature, and severity of AEs/SAEs/AESI, changes in laboratory data, vital signs, and ECGs compared with baseline, incidence and nature of study drug discontinuation, dose reduction, and dose delay due to AEs
 2. Efficacy - DoR and CR, ORR and CR rates
 3. Efficacy - DoCR, TTR, EFS, PFS, TTNT, OS
 4. PK - PK parameters such as Cmax, Tmax, Ctrough of surovatamig (at predefined intervals)
 5. Immunogenicity -pre-existing and induced ADA for surovatamig
 6. PROs - difference in severity levels in fatigue, pain, cognition and overall side-effect burden
 7. MRD-negative CR rate
-

Momentos temporales de evaluación secundaria

No aportado

Criterios de Inclusión

1. Participant must be 18 years old and above at the time of signing the ICF.
 2. ECOG performance status of 0 to 2
 3. Relapsed or refractory disease after at least 2 prior lines of systemic therapy
 4. Histologically confirmed diagnosis of B-NHL, based on local pathology report, as determined by WHO 2022 classification
 5. FDG-avid and measurable disease
 6. Adequate liver, hematological, renal and cardiac function.
- The above is a summary, other inclusion criteria details may apply

Criterios de Exclusión

1. Diagnosis of chronic lymphocytic leukemia, Burkitt lymphoma, or Richters transformation
 2. Active CNS involvement by B-NHL.
 3. Leukemic presentation of disease
 4. Prior neurotoxicity/ICANS or CRS
 5. History or presence of clinically relevant CNS pathology (based on investigator assessment)
 6. Prior allogeneic HSCT or solid organ transplantation within 24 weeks of first planned dose of surovatamig
- The above is a summary, other exclusion criteria details may apply

Calendario

(Última actualización: 01/04/2026)

Autorización 18/02/2025	Inicio de Ensayo 09/05/2025	Inclusión Primer Paciente 31/12/2025	Interrumpido No aportado	Reiniciado No aportado
Fin de reclutamiento No aportado	Fin prematuro (España) No aportado	Fin prematuro (Global) No aportado	Fin del ensayo en España No aportado	Fin del ensayo global No aportado

Promotor

Astrazeneca AB Sweden

Astraallen GartunaKarlebyhus Byggnad 674

Contact Person

AstraZeneca Clinical Study Information Center

+18772409479

information.center@astrazeneca.com

Monetary support: N/A

Tipo de promotor: Comercial

Ceim

CEIm Hospital Universitari de Bellvitge

Feixa Llarga, s/n - Antiguo Módulo Banco de Santander 08907 Hospitalet de Llobregat

presidenciaceic@bellvitgehospital.cat

Centros

Activo (26/05/2025)

Hospital Universitari Vall D Hebron

Barcelona

BARCELONA

Hematología

Activo (09/05/2025)

Hospital Universitario Fundacion Jimenez Diaz

Madrid

MADRID

Hematología

Activo (15/01/2026)

Hospital Universitario Quironsalud Madrid

Pozuelo De Alarcon

MADRID

Hematología

Medicamentos

AZD0486

SOLUTION FOR INFUSION

INTRAVENOUS INFUSION

-

Principios Activos: N/A

Experimental

AZD0486

SOLUTION FOR INFUSION

INTRAVENOUS INFUSION

-

Principios Activos: N/A

Experimental

Sin resultados

A trial to learn how well surovatamig (AZD0486) works and how safe it is in adults with certain blood cancers

State Recruiting	Type of participants Not provided	Age Ranges Older than 64 , Adults (18-64 years)
Gender Both	Phases Phase II	Expected Participants 188
Results No results	Low level of intervention No	Rare disease No
Geographic coverage Undetermined	Areas of the study treatment, safety, effectiveness, pharmacokinetics, pharmacodynamics, pharmacogenetics, pharmacogenomics	Advanced therapy NO

Information

Identifier 2023-505789-27-00 (CTIS)	Protocol Code No aportado
---	-------------------------------------

Therapeutic area
Diseases [C] - Hemic and Lymphatic Diseases [C15]

Investigated Disease
Relapsed or Refractory Non-Hodgkin Lymphoma

Scientific Title
A Modular Phase II, Single-arm, Multicenter, Open-label Study to Evaluate the Efficacy and Safety of Surovatamig (AZD0486) in Participants with Relapsed or Refractory B-cell Non-Hodgkin Lymphoma (SOUNDTRACK-B)

Rationale

Not provided

Main Objective

To evaluate the efficacy of surovatamig (AZD0486) single therapy administered at the Recommended Phase 2 Dose (RP2D) in relapsed or refractory B-cell NHL

Primary Endpoints

ORR, defined as the proportion of participants achieving either a PR or CR based on Lugano 2014 Response Criteria as determined by an IRC

Temporary moments of secondary assessment

Not provided

Secondary Objective

1. To assess the safety and tolerability of surovatamig
 2. To further evaluate the efficacy of surovatamig by assessment of DoR and CR rate
 3. To evaluate ORR and CR rate
 4. To evaluate the efficacy of surovatamig by evaluation of additional efficacy measures based on Lugano 2014 Response Criteria
 5. To characterize the PK of surovatamig
 6. To describe the incidence of immunogenicity
 7. To evaluate patient-reported tolerability of surovatamig, including severity of key treatment-related symptoms and overall side-effect burden
 8. To evaluate patient-reported severity of key disease-related symptoms, as well as the impact of disease on lymphoma-specific concerns and general HRQoL, while on surovatamig
 9. Evaluate the impact of surovatamig on MRD-negative CR rate
-

Secondary Endpoints

1. Safety and tolerability of surovatamig - incidence, nature, and severity of AEs/SAEs/AESI, changes in laboratory data, vital signs, and ECGs compared with baseline, incidence and nature of study drug discontinuation, dose reduction, and dose delay due to AEs
 2. Efficacy - DoR and CR, ORR and CR rates
 3. Efficacy - DoCR, TTR, EFS, PFS, TTNT, OS
 4. PK - PK parameters such as Cmax, Tmax, Ctrough of surovatamig (at predefined intervals)
 5. Immunogenicity -pre-existing and induced ADA for surovatamig
 6. PROs - difference in severity levels in fatigue, pain, cognition and overall side-effect burden
 7. MRD-negative CR rate
-

Temporary moments of secondary assessment

Not provided

Inclusion criteria

1. Participant must be 18 years old and above at the time of signing the ICF.
 2. ECOG performance status of 0 to 2
 3. Relapsed or refractory disease after at least 2 prior lines of systemic therapy
 4. Histologically confirmed diagnosis of B-NHL, based on local pathology report, as determined by WHO 2022 classification
 5. FDG-avid and measurable disease
 6. Adequate liver, hematological, renal and cardiac function.
- The above is a summary, other inclusion criteria details may apply

Exclusion criteria

1. Diagnosis of chronic lymphocytic leukemia, Burkitt lymphoma, or Richters transformation
 2. Active CNS involvement by B-NHL.
 3. Leukemic presentation of disease
 4. Prior neurotoxicity/ICANS or CRS
 5. History or presence of clinically relevant CNS pathology (based on investigator assessment)
 6. Prior allogeneic HSCT or solid organ transplantation within 24 weeks of first planned dose of surovatamig
- The above is a summary, other exclusion criteria details may apply

Calendar

(Last Update: 01/04/2026)

Authorization 18/02/2025	Start of Trial 09/05/2025	First patient inclusion 31/12/2025	Halted Not aported	Restarted Not aported
End of recruitment Not aported	Premature end (Spain) Not aported	Premature End (Global) Not aported	Trial end (Spain) Not aported	Trial end (Global) Not aported

Sponsor

Astrazeneca AB Sweden

Astraallen GartunaKarlebyhus Byggnad 674

Contact Person

AstraZeneca Clinical Study Information Center

+18772409479

information.center@astrazeneca.com

Monetary support: N/A

Sponsor type: Commercial

Ceim

CEIm Hospital Universitari de Bellvitge

Feixa Llarga, s/n - Antiguo Módulo Banco de Santander 08907 Hospitalet de Llobregat

presidenciaceic@bellvitgehospital.cat

Sites

Active (26/05/2025)	Hospital Universitari Vall D Hebron
	Barcelona BARCELONA Hematología
Active (09/05/2025)	Hospital Universitario Fundacion Jimenez Diaz
	Madrid MADRID Hematología
Active (15/01/2026)	Hospital Universitario Quironsalud Madrid
	Pozuelo De Alarcon MADRID Hematología

Medication

AZD0486 SOLUTION FOR INFUSION INTRAVENOUS INFUSION	-
Active Principles: N/A	Experimental
AZD0486 SOLUTION FOR INFUSION INTRAVENOUS INFUSION	-
Active Principles: N/A	Experimental

No results