

A Study to Investigate the Efficacy and Safety of Sonrotoclax Plus Zanubrutinib Compared With Placebo Plus Zanubrutinib in Adult Patients With Relapsed/Refractory Mantle Cell Lymphoma

Estado

Reclutando

Tipo de Participantes

No aportado

Rangos de Edad

Mayores de 64 , Adultos (18-64 años)

Género

Ambos

Fases

Fase III

Participantes esperados

160

Resultados

Sin resultados

Bajo nivel intervención

No

Enfermedad rara

Si

Cobertura geográfica

Sin determinar

Ámbitos del ensayo

tratamiento

Terapia avanzada

NO

Información

Identificador

2024-515593-27-00 (CTIS)

Cod. Protocolo

BGB-11417-302

Área terapéutica

Enfermedades [C] - Cáncer [C04]

Enfermedad investigada

relapsed or refractory mantle cell lymphoma (R/R MCL)

Título Científico

A Phase 3 Randomized Double-Blind Multicenter Study of Sonrotoclax Plus Zanubrutinib Versus Placebo Plus Zanubrutinib in Patients With Relapsed/Refractory Mantle Cell Lymphoma

Justificación

No aportado

Objetivo Principal

To demonstrate superiority of sonrotoclax plus zanubrutinib over placebo plus zanubrutinib as measured by progression-free survival (PFS) determined by a blinded independent review committee (BIRC).

Variables de Evaluación Primaria

Progression-free survival (PFS), as assessed by blinded independent review committee (BIRC), defined as the time from randomization to the date of progression or death, whichever occurs first.

Momentos temporales de evaluación primaria

No aportado

Objetivo Secundario

To compare the efficacy of sonrotoclax plus zanubrutinib with that of placebo plus zanubrutinib as measured by overall survival (OS).

To evaluate efficacy, as measured by the following: 1) Progression-free survival (PFS) determined by investigator. 2) Overall response rate (ORR), as determined by blinded independent review committee (BIRC) and by investigator assessment, per Lugano 2014 criteria. 3) Complete response rate (CRR), as determined by BIRC and by investigator. 4) Duration of response (DOR), as determined by BIRC and by investigator. 5) Time to first response, as determined by BIRC and by investigator.

To evaluate patient-reported disease and treatment symptoms.

To evaluate safety and tolerability.

Variables de Evaluación Secundaria

Overall survival (OS), defined as the time from randomization to the date of death from any cause.

Progression-free survival (PFS) as determined by investigator.

Overall response rate (ORR) as determined by BIRC and by investigator per the Lugano 2014 criteria. ORR is defined as the proportion of patients who achieved a best overall response of partial response or complete response (CR).

Duration of response (DOR) as determined by BIRC and by investigator. It is defined as the time from the first qualifying response to the date of progression or death, whichever occurs first.

Complete response rate (CRR), as determined by BIRC and by investigator. It is defined as the proportion of patients who achieved a best overall response of CR.

Time to first response as determined by BIRC and by investigator. It is defined as time from randomization to first response.

Time to initiation of new anticancer therapy.

Patient-reported symptom burden and physical condition/fatigue as measured by the European Organisation of Research and Treatment of Cancer-Quality of Life Questionnaire Non-Hodgkin Lymphoma High Grade Module 29 (EORTC-QLQ-NHL-HG29) questionnaire, and global health status (GHS) and physical function as measured by the European Organisation of Research and Treatment of Cancer-Quality of Life Questionnaire Core 30 (EORTC-QLQ-C30).

Incidence and severity of treatment-emergent adverse events (TEAE)s graded according to National Cancer Institute Common Terminology Criteria for Adverse Events (NCI-CTCAE) v5.0

Momentos temporales de evaluación secundaria

No aportado

Criterios de Inclusión

Histologically confirmed diagnosis of MCL based on the World Health Organization 2022 classification of Haematolymphoid Tumors (WHOHAEM5), or based on International Consensus Classification (ICC) Received 1 to 5 prior lines of systemic therapy including an anti-CD20 monoclonal antibody (mAb)-based immunotherapy or chemoimmunotherapy and requiring treatment in the opinion of the investigator Relapsed or refractory disease after the last line of therapy Measurable disease defined as 1 nodal lesion that is > 1.5 cm in longest diameter, or 1 extranodal lesion that is > 1 cm in longest diameter Eastern Cooperative Oncology Group (ECOG) Performance Status of 0 to 2 Adequate organ function

Criterios de Exclusión

Prior therapy with B-cell lymphoma-2 inhibitor
Prior therapy with covalent or non-covalent Bruton tyrosine kinase inhibitor (BTKi) unless the participant was intolerant of non-zanubrutinib covalent or non-covalent BTKi
Prior autologous stem cell transplantation or chimeric antigen receptor T-cell therapy within 3 months before first dose of study drug
Prior allogeneic stem cell transplant within 6 months of the first dose of the study drug
Known central nervous system involvement by lymphoma
Clinically significant cardiovascular disease
History of stroke or intracranial hemorrhage within 6 months before first dose of study drug

Calendario

(Última actualización: 06/07/2026)

Autorización 21/04/2025	Inicio de Ensayo 13/05/2025	Inclusión Primer Paciente 23/05/2025	Interrumpido No aportado	Reiniciado No aportado
Fin de reclutamiento No aportado	Fin prematuro (España) No aportado	Fin prematuro (Global) No aportado	Fin del ensayo en España No aportado	Fin del ensayo global No aportado

Promotor

BeOne Medicines AG Switzerland

Aeschengraben 27

Contact Person

BeOne Medical Officer

+18778285568

clinicaltrials@beonemed.com

Monetary support: N/A

Tipo de promotor: Comercial

Ceim

CEIm Regional de la Comunidad de Madrid

C/ Aduana 29 3ª planta 28013 Madrid

comite.regional@salud.madrid.org

913702824

Centros

Hospital Clinic De Barcelona

Barcelona

BARCELONA

Hematology

Activo (13/05/2025)

Hospital Universitario Fundacion Jimenez Diaz

Madrid

MADRID

Hematology

Activo (28/05/2025)

Hospital Universitario Puerta De Hierro De Majadahonda

Majadahonda

MADRID

Hematology

Activo (14/05/2025)

Institut Catala D'oncologia

L'hospitalet De Llobregat

BARCELONA

Hematology

Activo (16/07/2025)

Medicamentos

Zanubrutinib

CAPSULE
ORAL

-

Principios Activos: N/A

Experimental

BGB-11417

FILM-COATED TABLET
ORAL

-

Principios Activos: N/A

Experimental

BGB-11417

FILM-COATED TABLET
ORAL

-

Principios Activos: N/A

Experimental

BGB-11417

FILM-COATED TABLET
ORAL

-

Principios Activos: N/A

Experimental

BGB-11417

FILM-COATED TABLET
ORAL

-

Principios Activos: N/A

Experimental

Sin resultados

A Study to Investigate the Efficacy and Safety of Sonrotoclax Plus Zanubrutinib Compared With Placebo Plus Zanubrutinib in Adult Patients With Relapsed/Refractory Mantle Cell Lymphoma

State
Recruiting

Type of participants
Not provided

Age Ranges
Older than 64 , Adults (18-64 years)

Gender
Both

Phases
Phase III

Expected Participants
160

Results
No results

Low level of intervention
No

Rare disease
Yes

Geographic coverage
Undetermined

Areas of the study
treatment

Advanced therapy
NO

Information

Identifier
2024-515593-27-00 (CTIS)

Protocol Code
BGB-11417-302

Therapeutic area
Diseases [C] - Cancer [C04]

Investigated Disease
relapsed or refractory mantle cell lymphoma (R/R MCL)

Scientific Title
A Phase 3 Randomized Double-Blind Multicenter Study of Sonrotoclax Plus Zanubrutinib Versus Placebo Plus Zanubrutinib in Patients With Relapsed/Refractory Mantle Cell Lymphoma

Rationale

Not provided

Main Objective

To demonstrate superiority of sonrotoclax plus zanubrutinib over placebo plus zanubrutinib as measured by progression-free survival (PFS) determined by a blinded independent review committee (BIRC).

Primary Endpoints

Progression-free survival (PFS), as assessed by blinded independent review committee (BIRC), defined as the time from randomization to the date of progression or death, whichever occurs first.

Temporary moments of secondary assessment

Not provided

Secondary Objective

To compare the efficacy of sonrotoclax plus zanubrutinib with that of placebo plus zanubrutinib as measured by overall survival (OS).

To evaluate efficacy, as measured by the following: 1) Progression-free survival (PFS) determined by investigator. 2) Overall response rate (ORR), as determined by blinded independent review committee (BIRC) and by investigator assessment, per Lugano 2014 criteria. 3) Complete response rate (CRR), as determined by BIRC and by investigator. 4) Duration of response (DOR), as determined by BIRC and by investigator. 5) Time to first response, as determined by BIRC and by investigator.

To evaluate patient-reported disease and treatment symptoms.

To evaluate safety and tolerability.

Secondary Endpoints

Overall survival (OS), defined as the time from randomization to the date of death from any cause.

Progression-free survival (PFS) as determined by investigator.

Overall response rate (ORR) as determined by BIRC and by investigator per the Lugano 2014 criteria. ORR is defined as the proportion of patients who achieved a best overall response of partial response or complete response (CR).

Duration of response (DOR) as determined by BIRC and by investigator. It is defined as the time from the first qualifying response to the date of progression or death, whichever occurs first.

Complete response rate (CRR), as determined by BIRC and by investigator. It is defined as the proportion of patients who achieved a best overall response of CR.

Time to first response as determined by BIRC and by investigator. It is defined as time from randomization to first response.

Time to initiation of new anticancer therapy.

Patient-reported symptom burden and physical condition/fatigue as measured by the European Organisation of Research and Treatment of Cancer-Quality of Life Questionnaire Non-Hodgkin Lymphoma High Grade Module 29 (EORTC-QLQ-NHL-HG29) questionnaire, and global health status (GHS) and physical function as measured by the European Organisation of Research and Treatment of Cancer-Quality of Life Questionnaire Core 30 (EORTC-QLQ-C30).

Incidence and severity of treatment-emergent adverse events (TEAE)s graded according to National Cancer Institute Common Terminology Criteria for Adverse Events (NCI-CTCAE) v5.0

Temporary moments of secondary assessment

Not provided

Inclusion criteria

Histologically confirmed diagnosis of MCL based on the World Health Organization 2022 classification of Haematolymphoid Tumors (WHOHAEM5), or based on International Consensus Classification (ICC) Received 1 to 5 prior lines of systemic therapy including an anti-CD20 monoclonal antibody (mAb)-based immunotherapy or chemoimmunotherapy and requiring treatment in the opinion of the investigator Relapsed or refractory disease after the last line of therapy Measurable disease defined as 1 nodal lesion that is > 1.5 cm in longest diameter, or 1 extranodal lesion that is > 1 cm in longest diameter Eastern Cooperative Oncology Group (ECOG) Performance Status of 0 to 2 Adequate organ function

Exclusion criteria

Prior therapy with B-cell lymphoma-2 inhibitor
Prior therapy with covalent or non-covalent Bruton tyrosine kinase inhibitor (BTKi) unless the participant was intolerant of non-zanubrutinib covalent or non-covalent BTKi
Prior autologous stem cell transplantation or chimeric antigen receptor T-cell therapy within 3 months before first dose of study drug
Prior allogeneic stem cell transplant within 6 months of the first dose of the study drug
Known central nervous system involvement by lymphoma
Clinically significant cardiovascular disease
History of stroke or intracranial hemorrhage within 6 months before first dose of study drug

Calendar

(Last Update: 06/07/2026)

Authorization 21/04/2025	Start of Trial 13/05/2025	First patient inclusion 23/05/2025	Halted Not aported	Restarted Not aported
End of recruitment Not aported	Premature end (Spain) Not aported	Premature End (Global) Not aported	Trial end (Spain) Not aported	Trial end (Global) Not aported

Sponsor

BeOne Medicines AG Switzerland
Aeschengraben 27

Contact Person

BeOne Medical Officer
+18778285568
clinicaltrials@beonemed.com

Monetary support: N/A
Sponsor type: Commercial

Ceim

CEIm Regional de la Comunidad de Madrid
C/ Aduana 29 3ª planta 28013 Madrid
comite.regional@salud.madrid.org
913702824

Sites

Active (13/05/2025)	Hospital Clinic De Barcelona Barcelona BARCELONA Hematology
Active (28/05/2025)	Hospital Universitario Fundacion Jimenez Diaz Madrid MADRID Hematology
Active (14/05/2025)	Hospital Universitario Puerta De Hierro De Majadahonda Majadahonda MADRID Hematology
Active (16/07/2025)	Institut Catala D'oncologia L'hospitalet De Llobregat BARCELONA Hematology

Medication

Zanubrutinib

CAPSULE

ORAL

-

Active Principles: N/A

Experimental

BGB-11417

FILM-COATED TABLET

ORAL

-

Active Principles: N/A

Experimental

BGB-11417

FILM-COATED TABLET

ORAL

-

Active Principles: N/A

Experimental

BGB-11417

FILM-COATED TABLET

ORAL

-

Active Principles: N/A

Experimental

BGB-11417

FILM-COATED TABLET

ORAL

-

Active Principles: N/A

Experimental

No results