

A Phase 2 Multicohort Trial to Further Characterize the Efficacy and Safety of Ciltacabtagene Autoleucel

Estado

Reclutando

Tipo de Participantes

No aportado

Rangos de Edad

Mayores de 64 , Adultos (18-64 años)

Género

Ambos

Fases

Fase II

Participantes esperados

32

Resultados

Sin resultados

Bajo nivel intervención

No

Enfermedad rara

Si

Cobertura geográfica

Sin determinar

Ámbitos del ensayo

seguridad, eficacia, farmacocinética, dosis

Terapia avanzada

Terapia génica

Información

Identificador

2025-521975-30-00 (CTIS)

Cod. Protocolo

68284528MMY2012

Área terapéutica

Enfermedades [C] - Cáncer [C04]

Enfermedad investigada

Newly Diagnosed Multiple Myeloma

Título Científico

A Phase 2 Multicohort Trial to Further Characterize the Efficacy and Safety of Ciltacabtagene Autoleucel

Justificación

No aportado

Objetivo Principal

To characterize the efficacy of ciltacicel infusion following a fludarabine free LD regimen (Cohort A) or a low-dose ciltacicel infusion following a standard cyclophosphamide and fludarabine LD regimen (Cohort B)

Variables de Evaluación Primaria

MRD-negative CR at 12 months after ciltacicel infusion with a sensitivity of 10^{-5}

Momentos temporales de evaluación primaria

No aportado

Objetivo Secundario

No aportado

Variables de Evaluación Secundaria

No aportado

Momentos temporales de evaluación secundaria

No aportado

Criterios de Inclusión

1. At the time of informed consent, be 18 years of age (or at least the legal age of consent in the jurisdiction in which the study is taking place) but 80 years of age.
 2. Documented diagnosis of NDMM according to the most recent IMWG diagnostic criteria and measurable disease at diagnosis (prior to start of any anti-myeloma therapy)
 3. Not considered a candidate for high-dose chemotherapy with stem cell transplantation due to: -Advanced age; or - Presence of comorbid condition(s) likely to have a negative impact on tolerability of high-dose chemotherapy with stem cell transplantation; or -Participant refusal of high-dose chemotherapy with stem cell transplantation as initial treatment.
-

4. Participant must have received induction therapy. Initially, only participants receiving triplet induction therapy with DRd or VRd will be enrolled. Only after sponsor notification, participants receiving quadruplet DVRd induction therapy may be enrolled. Participants must have achieved PR to be enrolled on the most recent disease assessment.
5. ECOG Performance Status score of 0 or 1
6. Clinical laboratory values as defined in the protocol

Criterios de Exclusión

1. Prior systemic therapy for multiple myeloma or smoldering myeloma other than VRd, DRd or DVRd as per inclusion 4a and short course of corticosteroids . For participants who require emergency treatment, refer to protocol Section 4.4. 2. Radiation therapy for treatment of plasmacytoma within 14 days before enrollment. Palliative radiation for pain control secondary to lytic lesion is allowed within 14 days of enrollment, if the radiation portal covered 5% of the bone marrow reserve 3. Frailty index of 2 according to Myeloma Geriatric Assessment score 4. The following cardiac conditions: - New York Heart Association stage III or IV congestive heart failure. - Myocardial infarction or coronary artery bypass graft 6 months prior to enrollment. - History of clinically significant ventricular arrhythmia or unexplained syncope, not believed to be vasovagal in nature or due to dehydration. - History of severe non-ischemic cardiomyopathy. - Impaired cardiac function (LVEF <45%) as assessed by echocardiogram or multiple-gated acquisition scan (performed 60 days prior to apheresis). 5. Grade 2 or higher ongoing non-hematologic toxicity due to induction therapy, with the exception of grade 2 peripheral neuropathy due to bortezomib. 6. Participants who require continuous supplemental oxygen.

Calendario

(Última actualización: 24/07/2026)

Autorización 20/08/2025	Inicio de Ensayo 03/10/2025	Inclusión Primer Paciente 03/10/2025	Interrumpido No aportado	Reiniciado No aportado
-----------------------------------	---------------------------------------	---	------------------------------------	----------------------------------

Fin de reclutamiento No aportado	Fin prematuro (España) No aportado	Fin prematuro (Global) No aportado	Fin del ensayo en España No aportado	Fin del ensayo global No aportado
--	--	--	--	---

Promotor

Janssen Cilag International Belgium

Turnhoutseweg 30

Contact Person

CTIS Point of Contact

+31717996133

CTISadmin@its.jnj.com

Monetary support: N/A

Tipo de promotor: Comercial

Ceim

CEIm de Cantabria

Edificio IFIMAV 3ª Planta Avda Cardenal Herrera Oria s/n 39011 Santander

eclinicos4@idival.org

942315514

Centros

Activo (01/09/2025)	Clinica Universidad De Navarra Pamplona NAVARRA Hematologia
Activo (29/08/2025)	Hospital Clinic De Barcelona Barcelona BARCELONA Hematologia
Activo (23/01/2026)	Hospital Germans Trias I Pujol Badalona BARCELONA Hematologia
Activo (25/02/2026)	Hospital Universitari Vall D Hebron Barcelona BARCELONA Hematologia
Activo (28/08/2025)	Hospital Universitario De Salamanca Salamanca SALAMANCA Hematologia
Activo (30/09/2025)	Hospital Universitario Marques De Valdecilla Santander CANTABRIA Hematologia
Activo (27/08/2025)	Hospital Universitario 12 De Octubre Madrid MADRID Hematologia
Activo (30/09/2025)	University Hospital Virgen Del Rocio S.L. Sevilla SEVILLA Hematologia

Medicamentos

JNJ-68284528 DISPERSION FOR INFUSION INTRAVENOUS USE - Principios Activos: N/A Huérfano Experimental	CARVYKTI 3.2 &times; 10⁶ 1 &times; 10⁸ cells dispersion for infusion DISPERSION FOR INFUSION INTRAVENOUS USE Código ATC: L01XL05 - CILTACABTAGEN AUTOLEUCEL Principios Activos: N/A Huérfano Experimental
---	--

Sin resultados

A Phase 2 Multicohort Trial to Further Characterize the Efficacy and Safety of Ciltacabtagene Autoleucl

State
Recruiting

Type of participants
Not provided

Age Ranges
Older than 64 , Adults (18-64 years)

Gender
Both

Phases
Phase II

Expected Participants
32

Results
No results

Low level of intervention
No

Rare disease
Yes

Geographic coverage
Undetermined

Areas of the study
safety, effectiveness, pharmacokinetics, dose

Advanced therapy
Gene therapy

Information

Identifier
2025-521975-30-00 (CTIS)

Protocol Code
68284528MMY2012

Therapeutic area
Diseases [C] - Cancer [C04]

Investigated Disease
Newly Diagnosed Multiple Myeloma

Scientific Title
A Phase 2 Multicohort Trial to Further Characterize the Efficacy and Safety of Ciltacabtagene Autoleucl

Rationale

Not provided

Main Objective

To characterize the efficacy of ciltacabine infusion following a fludarabine free LD regimen (Cohort A) or a low-dose ciltacabine infusion following a standard cyclophosphamide and fludarabine LD regimen (Cohort B)

Primary Endpoints

MRD-negative CR at 12 months after ciltacabine infusion with a sensitivity of 10^{-5}

Temporary moments of secondary assessment

Not provided

Secondary Objective

Not provided

Secondary Endpoints

Not provided

Temporary moments of secondary assessment

Not provided

Inclusion criteria

1. At the time of informed consent, be 18 years of age (or at least the legal age of consent in the jurisdiction in which the study is taking place) but 80 years of age.
 2. Documented diagnosis of NDMM according to the most recent IMWG diagnostic criteria and measurable disease at diagnosis (prior to start of any anti-multiple myeloma therapy)
 3. Not considered a candidate for high-dose chemotherapy with stem cell transplantation due to: -Advanced age; or - Presence of comorbid condition(s) likely to have a negative impact on tolerability of high-dose chemotherapy with stem cell transplantation; or -Participant refusal of high-dose chemotherapy with stem cell transplantation as initial treatment.
-

4. Participant must have received induction therapy. Initially, only participants receiving triplet induction therapy with DRd or VRd will be enrolled. Only after sponsor notification, participants receiving quadruplet DVRd induction therapy may be enrolled. Participants must have achieved PR to be enrolled on the most recent disease assessment.
5. ECOG Performance Status score of 0 or 1
6. Clinical laboratory values as defined in the protocol

Exclusion criteria

1. Prior systemic therapy for multiple myeloma or smoldering myeloma other than VRd, DRd or DVRd as per inclusion 4a and short course of corticosteroids . For participants who require emergency treatment, refer to protocol Section 4.4. 2. Radiation therapy for treatment of plasmacytoma within 14 days before enrollment. Palliative radiation for pain control secondary to lytic lesion is allowed within 14 days of enrollment, if the radiation portal covered 5% of the bone marrow reserve 3. Frailty index of 2 according to Myeloma Geriatric Assessment score 4. The following cardiac conditions: - New York Heart Association stage III or IV congestive heart failure. - Myocardial infarction or coronary artery bypass graft 6 months prior to enrollment. - History of clinically significant ventricular arrhythmia or unexplained syncope, not believed to be vasovagal in nature or due to dehydration. - History of severe non-ischemic cardiomyopathy. - Impaired cardiac function (LVEF <45%) as assessed by echocardiogram or multiple-gated acquisition scan (performed 60 days prior to apheresis). 5. Grade 2 or higher ongoing non-hematologic toxicity due to induction therapy, with the exception of grade 2 peripheral neuropathy due to bortezomib. 6. Participants who require continuous supplemental oxygen.

Calendar

(Last Update: 24/07/2026)

Authorization 20/08/2025	Start of Trial 03/10/2025	First patient inclusion 03/10/2025	Halted Not aported	Restarted Not aported
------------------------------------	-------------------------------------	--	------------------------------	---------------------------------

End of recruitment Not aported	Premature end (Spain) Not aported	Premature End (Global) Not aported	Trial end (Spain) Not aported	Trial end (Global) Not aported
--	---	--	---	--

Sponsor

Janssen Cilag International Belgium

Turnhoutseweg 30

Contact Person

CTIS Point of Contact

+31717996133

CTISadmin@its.jnj.com

Monetary support: N/A

Sponsor type: Commercial

Ceim

CEIm de Cantabria

Edificio IFIMAV 3ª Planta Avda Cardenal Herrera Oria s/n 39011 Santander

eclinicos4@idival.org

942315514

Sites

Active (01/09/2025)	Clinica Universidad De Navarra Pamplona NAVARRA Hematologia
Active (29/08/2025)	Hospital Clinic De Barcelona Barcelona BARCELONA Hematologia
Active (23/01/2026)	Hospital Germans Trias I Pujol Badalona BARCELONA Hematologia
Active (25/02/2026)	Hospital Universitari Vall D Hebron Barcelona BARCELONA Hematologia
Active (28/08/2025)	Hospital Universitario De Salamanca Salamanca SALAMANCA Hematologia
Active (30/09/2025)	Hospital Universitario Marques De Valdecilla Santander CANTABRIA Hematologia
Active (27/08/2025)	Hospital Universitario 12 De Octubre Madrid MADRID Hematologia
Active (30/09/2025)	University Hospital Virgen Del Rocio S.L. Sevilla SEVILLA Hematologia

Medication

JNJ-68284528 DISPERSION FOR INFUSION INTRAVENOUS USE - Active Principles: N/A Orphan Experimental	CARVYKTI 3.2 $\times 10^6$ $\times 10^8$ cells dispersion for infusion DISPERSION FOR INFUSION INTRAVENOUS USE ATC code: L01XL05 - CILTACABTAGEN AUTOLEUCEL Active Principles: N/A Orphan Experimental
--	---

No results